

Surgery ~ DO NOT TAKE MEDICATION LIST

The following is a list of medications that cause thinning of the blood. These medications should be avoided 2 weeks BEFORE & AFTER SURGERY secondary to the risk of excessive bleeding during and / or after surgery. YOU MAY TAKE TYLENOL FOR PAIN DURING THIS 4 WEEK PERIOD.

Advil	Clinoril	Ketorlac
Aggrenox	Coumadin	Ketoprofen
Aleve	Darvon	Lodine
Alka Seltzer	Daypro	Magsal
Amigesic Capsules	DiFlunisal	Marnal
Anacin	Doan's	Marthritic
Anaprox	Dolobid	MecloFenamate
Ansaid	Dristan	Meclomen
Argesic	Easprin	Menadol
Arthritis Bayer Timed Release	Ecotrin	Micrainin
Arthritis Pain	Effient	Midol
Arthritis Strength Bufferin	Eliquis	Mobidin
Arthropan	Emagrin	Mobigesic
Arthrotec	Empirin	Motrin
Ascriptin	Endodan	Nalfon
Aspergum	Equageic	Naprosyn
Aspercreme	Etodolac	Norgesic
Aspirin or Aspirin Products	Excedrin	Norwich
Axotal	Feldene	Nuprin
Azulfidine (Sulfasalazine)	Fenoprofen	Orphengesic
Bayer Aspirin	Fiornal	Orudis
BC Powder	Fluriprofen	Oruvail
Ben Gay	Goody's	PAC Tablets
Brufen	Halfprin	Pamprin
Bufferin	Haltram	Pesto-Bismol
Butazolidin	Ibuprin	Percodan
Cama Arthritis Pain Reliever	Ibuprofen	Piroxicam
Cataflam	Indocin	Plavix
Cheracol	Indomethacin	Ponstel

Pradaxa	St. Joseph Aspirin	Treximet
Relafen	Sulindac	Tricosal
Furen	Supac	Triilisate
Saleto Products	Synalgos	Vanquish
Salflex	Talwin	Voltaren
Salsalate	Ticlid	Warfarin
Salsitabs	Tolectin	Xarelto
SineAid	Tolmectin	Zorpin
SineOff	Toradol	
Soma	Trendar	

ALL HERBAL SUPPLEMENTS

Ginko, Black Cohosh, Juice Plus, Green Tea, Red Wine

ALL DIET MEDICATIONS - PRESCRIPTION & OVER-THE-COUNTER

VITAMINS ABOVE THE RECOMMENDED DAILY ALLOWANCE